

---

# FOCUS: MARINE GEOMORPHOLOGY AS A DETERMINANT FOR ESSENTIAL LIFE HABITAT AND MARINE PROTECTED AREA DESIGN

---

## Marine Geomorphology in the Design of Marine Reserve Networks\*

**William D. Heyman**

*Texas A&M University, College Station*

**Dawn J. Wright**

*Oregon State University*

Marine environments, key life-support systems for the earth, are under severe threat. Issues associated with managing these common property resources are complex and interrelated. Networks of marine reserves can be valuable for mitigating threats to marine systems, yet the successful design and implementation of such networks has been limited. Efficient ways to conserve marine environments are urgently needed. This Focus Section of *The Professional Geographer* explores the development of marine reserve networks based on geomorphology, fish biology, ecological connectivity, and appropriate governance. The articles in this Focus Section offer examples of the following: (1) distinctive reef geomorphology dictating the spawning locations of reef fishes, which in turn serve as critical source sites for the replenishment of distant reefs by means of larval transport; (2) an example of a simplified oceanographic model that predicts larval transport from fish breeding sites to important nursery areas; and (3) a case study of the development of a marine reserve network that illustrates key elements of a successful strategy. In sum, this Focus Section offers case studies that show the value of marine geomorphology, oceanographic connectivity, and stakeholder involvement as key elements of multidisciplinary geographic studies applied to the design of marine reserve networks. Geographers can further contribute to the conservation and management of coastal and marine ecosystems in many ways that involve subdisciplines of remote sensing and geographic information systems, political and economic geography, political ecology, and ethnography. **Key Words: connectivity, geomorphology, governance of common property resources, marine reserves, spatial planning, spawning.**

海洋环境是地球生命支持的关键系统，正在受到严重威胁。与这些共同财产资源管理有关的问题是复杂和相互关联的。海洋保护区网络对减轻海洋系统的威胁是很可贵的，但目前对这种网络的成功设计和实施很有限。我们迫切需要能够保护海洋环境的有效方法。《专业地理学家》的重点章节探讨了以地貌、鱼类生物学、生态连接、和适度管理为基础的海洋保护区网络的发展。上述重点章节的文章提供了下列范例：（1）独特的珊瑚礁地貌支配了珊瑚鱼的产卵地点，这反过来通过幼虫运输，又成为了补给远方珊瑚礁的重要来源地点；（2）以一个简单的海洋地理模型为例，预测了鱼类幼虫从滋生地到重要育苗区的运输；以及（3）以一个海洋保护区网络的

\*We are grateful for all of the participation in two symposia at AAG meetings in 2008 and 2009 that led to the development of this special Focus Section. We are also grateful to the authors of this Focus Section for their hard work and continued efforts. Finally, we are grateful to the editor and staff of *The Professional Geographer*, without whom this Focus section could not have come to pass.

发展为个例，展示了一个成功策略所需的关键要素。总之，这个重点章节所提供的案例研究展示了海洋地貌学、海洋地理连接性、和利益相关者参与的价值，这些都是设计海洋保护区网络所需要应用的多学科地理研究的要素。借助于涉及多学科遥感地理信息系统、政治和经济地理学、政治生态学、和人种学，地理学家们可以进一步促进沿海和海洋生态系统的保护和管理。关键词：连通，地貌，共同财产资源管理，海洋保护区，空间规划，产卵。

Los entornos marinos, sistemas de apoyo biológico claves para la tierra, se encuentran severamente amenazados. Los asuntos asociados con el manejo de estos recursos de propiedad común son complejos e interrelacionados. Las redes de reservas marinas pueden ser valiosas para mitigar las amenazas a los sistemas marinos, aunque el diseño e implementación exitosos de tales redes ha sido de poco alcance. Se necesitan con urgencia procedimientos eficientes para conservar el medio ambiente marino. Esta Sección Focal de *The Professional Geographer* explora el desarrollo de ese tipo de redes en lo que concierne a geomorfología, biología ictiológica, conectividad ecológica y apropiada acción gubernamental. Los artículos de esta Sección Focal ofrecen ejemplos de lo siguiente: (1) la peculiar geomorfología coralina que determina las localidades de desove de los peces de arrecife, que a la vez sirven de asiento a fuentes críticas para el reabastecimiento de arrecifes distantes a través del transporte de larvas; (2) un ejemplo de un modelo oceanográfico simplificado que predice el transporte de larvas desde los sitios de reproducción de peces hasta áreas importantes de crecimiento; y (3) un estudio de caso del desarrollo de una red de reserva marina que ilustra sobre los elementos claves de una estrategia exitosa. En suma, esta Sección Focal ofrece estudios de casos que muestran el valor de la geomorfología marina, la conectividad oceanográfica y la activa participación de partes interesadas, como elementos claves de estudios geográficos multidisciplinarios aplicados al diseño de redes de reservas marinas. Los geógrafos pueden contribuir todavía más en la conservación y manejo de ecosistemas marinos litorales de maneras variadas que involucren las subdisciplinas de percepción remota y sistemas de información geográfica, geografía política y económica, ecología política y etnografía. **Palabras clave:** conectividad, geomorfología, administración de recursos de propiedad común, reservas marinas, planificación espacial, desove marino.

Marine waters cover 71 percent of the Earth's surface. They serve a key role in controlling Earth's climate and supporting human economies and social welfare. The annual value of coastal and marine ecosystem services was estimated at \$22.6 trillion, more than double that of terrestrial ecosystem services, \$10.7 trillion, and even greater than global domestic product which was estimated at \$18 trillion (Costanza et al. 1997). Nearly half of the world's population lives within 200 km of a coastline and that figure is likely to double by 2025 (Creel 2003). As the receiving basin for runoff and pollution and the last true commons left on Earth, marine waters are being degraded due to ocean acidification, overfishing, pollution, and habitat destruction; yet they continue to be managed poorly (Millennium Ecosystem Assessment 2005; Hoegh-Guldberg et al. 2007). The human and financial resources available for marine conservation and management are estimated as two orders of magnitude lower than required (Balmford et al. 2004). Understandably, efficient ways to conserve marine environments are urgently needed and have been the focus of increasing scientific and political attention.

It has been ten years since *The Professional Geographer* has addressed marine environments in a Focus Section (Steinberg 1999). That section illustrated the wide and growing body of geographic studies on marine and coastal environments from physical, human, political, and economic perspectives. Geographers have increasingly focused on marine ecosystem dynamics and management during the last decade (e.g., St. Martin 2001; Burne and Parvey 2002; Psuty, Steinberg, and Wright 2004; Lunn and Dearden 2006; Prigent et al. 2008), yet these efforts are still limited compared to the disciplinary emphasis on terrestrial landscapes. Rather than attempt to review all new marine geographic studies, this article offers examples of the diversity of contributions that geographers have made to marine and coastal management and demonstrates ways in which geographers could offer a more holistic approach to this vast study area. The overall objective of the Focus Section is to promote improved planning for marine reserve networks through the use of geomorphologic habitat proxies, studies of ecological habitat connectivity, and the involvement of local fishers and their local knowledge to conserve reef fish spawning aggregations.

We organized four sessions at the 2008 Association of American Geographers (AAG) annual meeting under the common theme, "Marine Geomorphology as a Determinant for Essential Life Habitat: An Ecosystem Management Approach to Planning for Marine Reserve Networks" (Wright and Heyman 2008b). The sessions were cosponsored by three specialty groups of the AAG: Coastal and Marine, Geographic Information Science and Systems, and Biogeography. The unifying goal of these sessions was to examine critically the growing body of data suggesting that, even more than in terrestrial environments, the underlying geology and geomorphology of marine environments dictates the location of critical life habitats for a variety of species. The broad implications of these findings suggest that geomorphology might be used as a proxy for (or at least help to identify) critical life habitat for marine species and thus serve to advance the applications of ecosystem-based management (EBM), the design of marine reserve networks, and marine spatial planning more generally.

The intentions of this lead article of this Focus Section are twofold: to (1) engage a wide array of scholars about the values and condition of marine waters and ways in which geographers can further contribute to marine and coastal management and (2) provide the context for the articles in this section that together focus on the design of marine reserves based on principles of geomorphology, environmental biology of fishes, connectivity, and the involvement of stakeholders in governance. The article navigates relevant background literature to explain key terms, concepts, and themes. The scope is necessarily broad and includes sections on the status and trends in marine fisheries and ecosystems, biology of marine fishes, marine geomorphology as proxy for marine habitats, marine remote sensing and habitat mapping, marine EBM, connectivity and larval transport, fisher traditional ecological knowledge (TEK), and stakeholder involvement in and governance of common property resources. These disparate themes converge to address marine EBM and marine spatial planning and form a synthesis of the Focus Section. This article thus serves as an introduction to and synthesis of the articles in the Focus Section and a review of the wide-ranging and important roles that geographers play in marine conservation and management.

## **Status and Trends in Marine Fisheries and Ecosystems**

Seafood produced from marine fisheries and aquaculture provides about 15 percent of the protein consumed by humans, more than 50 percent in small island developing states, and is the world's most highly traded food internationally. Net exports of fish and fishery products were valued at US\$24.6 billion in 2006, representing 194 participating countries (Food and Agriculture Organization of the United Nations [FAO] 2009). Our global dependence on marine fisheries and associated marine ecosystems is not often considered for its important role in global food security (Smith et al. 2010). Nonetheless, marine fisheries resources and the habitats on which they depend are either fully exploited or in decline throughout the world (National Research Council 1999; FAO 2004, 2009; U.S. Commission on Ocean Policy 2004; Millennium Ecosystem Assessment 2005). Low-latitude areas (e.g., the Caribbean) that harbor coral reef environments and a high proportion of the world's biodiversity exhibit rapid declining trends that are consistent with global averages (Burke and Maidens 2004). These low-latitude areas also have a high percentage of the world's poor, a higher percentage of people that are directly dependent on marine resources for protein and livelihoods, and often less effective governance structures (FAO 2009). Three case studies in this issue (Heyman; Coleman, Scanlon, and Koenig; Gleason, Kellison, and Reid) focus on the Gulf and Caribbean region, providing a look at low-latitude areas that have a variety of governance arrangements. The final study in this section (Fischer et al.) is from the California coast where resources for governance are more plentiful, although the study proposes a cost-effective way to go about marine conservation planning that could be applied in other areas.

There is strong and growing evidence that industrial fisheries are, by nature, unsustainable and have led to declines in marine and fishery resources, particularly large predators (Pauly et al. 2002; Myers and Worm 2003). There is also a growing realization that a variety of additional factors are affecting the health, resilience, biodiversity, and productivity of marine waters and the ocean's ability to produce the variety of ecosystem services on which societies depend

(Worm et al. 2006). Recreational fisheries, for example, can have enormous effects on fished stocks. Coleman et al. (2004) reported that 64 percent of the landings for species of concern in the Gulf of Mexico are harvested within recreational fisheries. This is particularly the case when recreational fishers target vulnerable times and places in fishes' life history (see the next section). Recreational fishers are numerous yet typically smaller producers than commercial fishers, and they have strong links to local tourism economies. As a result, their impacts and needs for careful management and regulation have been largely overlooked (Coleman et al. 2004).

Overfishing alters marine environments in a variety of ways. Jackson et al. (2001) have shown that overfishing over centuries has dramatically altered marine environments. They used historical data gleaned from paleoecological sedimentary records, archaeological records of human coastal settlements dating back 10,000 years, and historical records and charts dating back to the fifteenth century. Pauly et al. (1998) described "fishing down the food web" as the trophic consequences of overfishing whereby societal preference for large predatory species has created a serial top-down depletion that is having cascading effects throughout marine ecosystems. Global environmental changes also contribute directly to observed declines in marine ecosystem health and fisheries harvests. Hoegh-Guldberg et al. (2007) illustrated how rising ocean temperatures and acid concentrations have together contributed to the global decline in coral reef habitat extent and health. Riverborne sediments, nutrients, pesticides, and herbicides from upland agriculture, industry, and urban areas are also having major effects on coastal and marine ecosystems. The most deleterious effects are generally on near-shore habitats such as mangroves, seagrasses, and estuaries, which often serve as nursery grounds for a variety of species (Beck et al. 2001).

Management responses to the plethora of threats to marine systems are as varied as the problems. Unfortunately, sectoral, single-species, top-down approaches that have been imposed by fisheries regulators and management agencies have rarely proven effective. Worm et al. (2009) suggested that solutions to the global fisheries crisis must not focus sim-

ply on marine fisheries management interventions. Many authors have suggested a much more holistic approach to fisheries and marine management that is based on maintaining healthy and resilient marine ecosystems, recognizes connectivity, is spatially explicit, and is implemented through broad-sector participation at the largest possible scales (e.g., Crowder et al. 2008; Palumbi et al. 2009; Worm et al. 2009; Norse 2010; section on management of common property recurses later).

### **Biology of Marine Fishes as Relevant to Marine Management**

Distinct from terrestrial organisms, many marine fish species incorporate three periods of dispersal during their life history—a period of pelagic larval dispersal, ontogenetic habitat shifts through juvenile development, and seasonal adult migration for reproduction. Although there are some species that remain sedentary as adults and others that have very limited larval dispersal, nearly all fish release pelagic eggs (Claydon 2004). The persistence of each species and, by extension, the overall resilience of marine systems therefore depends on the availability of healthy areas for each life stage and successful movement or connectivity between them (Leis 1987; Roberts 1997; Peterson et al. 2000; Beck et al. 2001; Grober-Dunsmore and Keller 2008).

Most large-bodied, long-lived reef fishes do not spawn within their home range. Instead, they perform seasonal migrations for broadcast spawning from within transient aggregations to produce masses of pelagic larvae for dispersal (Claydon 2004). Fishes commonly migrate great distances to spawn within aggregations that occur at specific times and places (Johannes 1978; Thresher 1984; Leis 1987; Domeier and Colin 1997). Spawning aggregations of reef fish present easy targets for fishermen with unusually high densities of fishes at predictable times and areas (Johannes 1998; Sadovy De Mitcheson et al. 2008). Although some species appear more vulnerable to aggregation fishing than others, even light levels of fishing appear to affect the viability and health of spawning aggregations (Koenig et al. 1996; Sadovy and Domeier 2005). Fishing a species during its spawning aggregation has invariably led to declines and, in many cases, localized

extirpations (Johannes 1998; Sala, Starr, and Ballesteros 2001; Claro and Lindeman 2003; Sadovy De Mitcheson et al. 2008). Protecting reef fish spawning aggregations is an obvious conservation strategy (Johannes 1998) consistent with ecosystem-based fishery management (Pikitch et al. 2004; see later). Nonetheless, clear patterns of the timing and location of *multispecies* reef fish spawning aggregations are beginning to emerge. Several of the articles in this Focus Section examine explicitly the geomorphologically based marine habitats associated with spawning aggregations (Heyman; Gleason, Kellison, and Reid; Coleman, Scanlon, and Koenig).

### Marine Geomorphology as Proxy for Marine Habitats

Marine environments and their associated biota are dictated by their physical oceanographic and geographic setting at all scales. Classic geomorphological studies of landform have been eclipsed by more modern studies of process and dynamics (Psuty, Steinberg, and Wright 2004; Wright and Heyman 2008a). In marine environments, however, where bathymetric data are limited in scale and extent, landform (or submarine shape and form) studies are highly relevant and yet still somewhat rare. Yet geographic setting is fundamental in defining the structure and function of marine ecosystems. Coastal margin shapes (trailing vs. leading edge coasts), for example, are created by tectonic activity. Underlying geology provides the basis for the development of benthic habitats (e.g., sediment vs. rock). Water column properties such as temperature range, seasonal light variation, and tidal variation are functions of latitude. Species composition varies with hemisphere and region. The arrival and departure of regularly occurring but stochastic ocean gyres control local oceanic conditions. Far-field and localized winds influence wave height, period, and intensity, and each of these is attenuated by local structure. At smaller scales, biotic habitats provide structure for other species (e.g., coral reefs provide habitat for marine plants, invertebrates, and fishes). Indeed, coral reef habitats have been suggested as surrogates for species, ecological functions, and ecosystem services (Mumby et al. 2008).

Ecosystems consist of both biotic and abiotic components and their interactions. The diversity and density of species and their ecological relationships are generally difficult to observe and quantify particularly over large geographic areas, but communities of organisms are generally constrained by their physical environment. Marine biogeography probably began with the observations of Charles Darwin. More recently there have been various attempts to classify, characterize, and map marine environments at various scales (e.g., Hedgpeth 1957a, 1957b; Hayden, Ray and Dolan 1984; Lanier, Romsos, and Goldfinger 2007). A growing number of authors suggest that abiotic ecosystem attributes can be used as surrogates for the identification, mapping, and conservation of biotic components of ecosystems. This approach is fundamental to landscape ecology and, although challenging, is being increasingly adopted for marine systems (Pittman, McAlpine, and Pittman 2004; Pittman, Caldow, and Hile 2007; Grober-Dunsmore et al. 2008; Costello 2009).

Hierarchical classifications can be used to develop marine conservation strategies at regional, national, and global levels. Zacharias et al. (1998) offered an ecosystem classification scheme for British Columbia. Roff and Taylor (2000) provided an example of this approach used for the marine waters of Canada, the country with the longest coast and bordering three oceans. The hierarchical geophysical approach is supported by available data derived from remote sensing, bathymetric maps, and ocean circulation patterns (Zacharias and Roff 2000). Oceanographic and physiographic data are used to derive a consistent set of habitat classifications that together make up the seascape. Roff, Taylor, and Laughren (2003) argued that geophysical surrogates for marine community types are fundamental to understanding biotic distribution and thus the most practical foundation for marine planning, management, and conservation. In the same vein, geomorphology serves as a basis for a national conservation framework for the marine waters of Australia (Burne and Parvey 2002; Heap and Harris 2008). Global marine classifications following a similar geophysical approach are beginning to emerge as a basis for global marine planning and management (e.g., Spalding et al. 2007; Andréfouët et al. 2008).

A growing number of studies in a variety of locations are testing the validity of geophysical classification systems used to identify biological habitats (e.g., Wilson et al. 2007; Erdey-Heydorn 2008; Lampietro, Young, and Kvitek 2008; Kracker, Kendall, and McFall 2008; Wedding and Friedlander 2008). These articles support the concept with specific examples; they refine techniques and applications for marine and coastal planning, conservation, and marine reserve network design.

There is emerging evidence that many species that migrate to spawn aggregate at locations with particular geomorphic structures: generally abrupt discontinuities in surrounding structure such as reef promontories, uplifted ridges, and shelf edges. Several articles provided in this Focus Section (Gleason, Kellison, and Reid; Coleman, Scanlon, and Koenig; Heyman) provide evidence from east and west Florida and Belize, respectively, to support this claim. These patterns are further supported by the locations of reef fish spawning aggregations at similar geomorphological features in Cuba, the Cayman Islands, and other areas (Claro and Lindeman 2003; Whaylen et al. 2004; Kobara and Heyman 2008, 2010). Collisions between large-scale ocean currents (e.g., gyres) and abrupt changes in geomorphology alter localized oceanic conditions. These oceanographic and physical discontinuities create underlying ecosystem processes or conditions to which many species have been attracted over evolutionary time (Heyman and Kjerfve 2008). Together these serve as examples and provide evidence for the larger concept that geology and geomorphology must be taken into account in the design of EBM strategies.

### **Marine Remote-Sensing, Bathymetric Mapping, and Habitat Classifications**

In spite of the critical need for geomorphology and hence marine habitat information, the collective effort to map the seafloor has only produced accurate coverage for 5 to 10 percent of the world's seafloor (Sandwell et al. 2003; Wright 2003). Nonetheless, satellite- and aircraft-based remote-sensing techniques, ship-based single- and multibeam techniques, videography from free-swimming and towed diver surveys, remotely operated vehicles, sub-

mersibles, and computer-assisted geoprocessing advances have all contributed to a greater availability of marine habitat mapping techniques and products (as reviewed by Green et al. 1996; Wright 1999; Wright and Heyman 2008a). As a result, there have been dramatic increases in the extent and quality of marine geomorphological habitat characterizations and interpretations (e.g., Wright, Donahue, and Naar 2002; Aswani and Lauer 2006; Lanier, Romos, and Goldfinger 2007; Wilson et al. 2007; Kendall and Miller 2008).

In addition to habitat mapping, a discussion of geographically based marine reserve network designs would be incomplete without mention of the scores of geographic information system (GIS)-based spatial algorithms that have been developed for marine reserve planning and decision support (NatureServe 2008). One of the most notable is the suite of algorithms known as MARine reserve design using spatially eXplicit ANnealing (MARXAN; Ball and Possingham 2000; Possingham, Ball, and Andelman 2000; Leslie et al. 2003; Klein et al. 2008). MARXAN uses stochastic optimization routines to generate viable spatial reserve solutions that optimize coverage of preselected biological criteria, while minimizing the cost of the reserve network.

### **Marine Ecosystem-Based Management**

McLeod et al. (2005) defined EBM as an adaptive resource management approach that incorporates ecosystem processes and their responses to environmental perturbations while also addressing the complexity of human social systems (e.g., fishing communities, conservation organizations, local resource users, academic and research scientists, community members with traditional knowledge, and other stakeholders). EBM should increase the resilience of marine systems in the face of increasing local and global threats (Levin and Lubchenco 2008).

Marine EBM was once and still is common in a number of Pacific Island nations, where modern impacts have been relatively limited. These "traditional management" techniques have a great deal in common with what is presently being called EBM. Industrialized nations are

just now rediscovering these simple principles, which are particularly valuable for marine systems with high diversity. Coral reef ecosystems are high in diversity and their fisheries are concomitantly diverse; many species are targeted in small numbers. Reef fisheries are therefore difficult to manage with conventional, single-species management means such as quotas, size and bag limits, or closed seasons (Koenig et al. 2000). Instead, an ecosystem-based fishery management (EBFM) approach might be more effective (Pikitch et al. 2004). This approach, which differs slightly from EBM but is complementary, recognizes the interdependence between protection of critical life habitat and multispecies fishery production. Ecosystem functions and critical life habitat, such as spawning and nursery grounds, are protected from destructive fishing practices to promote sustainable harvests (Koenig et al. 2000; Pikitch et al. 2004). Recognizing that marine management includes more than just fishing, recent studies are urgently recommending broader EBM (e.g., McLeod et al. 2005; Christie et al. 2007; Crowder et al. 2008) to account for impacts on nontarget fisheries, resources, and habitats (e.g., nonpoint source pollution) and to include more of a participatory approach to management with broad stakeholder involvement. In part because both EBM and EBFM are new and complex in modern cultures, recent successful examples are uncommon (Crowder et al. 2008). Heyman (this issue) provides a case study from Belize where the broad participation of a diverse group of stakeholders (including local fisherman) and a detailed analysis of geomorphology and its association with the biology of exploited species play critical roles in the development of a national network of marine reserves. Indeed, most scientists agree in principle that large and functional marine reserve networks that provide connectivity between various critical habitats do form an essential (but not sufficient) component of any EBM or EBFM approach (e.g., National Research Council 1999, 2001; McLeod et al. 2005; Halpern et al. 2008; Norse 2010).

Marine reserves are therefore considered effective tools to manage fisheries and mitigate pressures on marine biodiversity (Roberts 1997; National Research Council 1999, 2001; Allison et al. 2003; Hastings and Botsford

2003). The optimal design of reserve networks has received a great deal of attention from modelers, ecologists, and managers, but generally the practical utility of these models and their outputs have been limited by the lack of biological data (e.g., distribution of species, larval behavior) with which to run the models (Roberts 1997; Halpern 2003; Halpern and Warner 2003; Hastings and Botsford 2003; Berkeley et al. 2004; U.S. Commission on Ocean Policy 2004).

There is an urgent need to rapidly expand the coverage of marine reserve networks to promote marine ecosystem management. Deciding where to place these reserves is a daunting task, particularly given political opposition and the paucity of available data on which to make decisions. Because bathymetric data have been primarily collected to assist navigation, the world's ports and large areas of shallow U.S. coastal waters have been extensively mapped. Beyond that, however, there exist astonishingly little fine-scale marine bathymetric data, especially for deep and remote areas. Sparsely available marine biogeographical data represent an even greater knowledge gap. We propose that geomorphological habitat proxies, based on the best available bathymetric data, can assist managers in making timely recommendations for inclusion of critical habitats within marine reserve networks. We urge, therefore, that bathymetric data collection and habitat mapping efforts be expanded to this end.

### **Connectivity and Larval Transport**

---

Maintaining "connectivity" between ecosystem components is critical for their effective maintenance, resilience, and survival and therefore must be considered in the design of marine reserve networks (Roberts 1997). Even if sufficient critical habitats are encompassed within the reserve network, managers are cognizant that most species immigrate and emigrate from reserves, both by swimming and by larval transport. Roberts et al. (2001) illustrated the positive "spillover effect" of reserves on adjacent fisheries. A variety of studies have addressed larval connectivity, generally through numerical circulation modeling. Warner, Swearer, and Caselle (2000) evaluated the relative importance of larval retention versus long-distance

transport of gametes for the design of marine reserve networks.

Fischer et al. (this issue) introduce an alternate approach, which focuses on larval exchange as a critical factor in marine reserve network design. They have developed a two-dimensional, GIS-based diffusion model for representing larval dispersal distributions based only on bathymetry and coastal oceanographic circulation patterns. The method holds great promise for practitioners attempting to design marine reserves with limited time and oceanographic information (i.e., limited access to complex particle-tracking models that might not even be available for a region in which a reserve network is being designed). The method is superior to standard one-dimensional approaches currently in use that estimate dispersal along a coastline in an advection-diffusion framework (e.g., Okubo and Levin 2002). Connectivity research is clearly an important area requiring further study and provides another valuable avenue for interested geographers with skills in biogeography and physical processes.

### Traditional Ecological Knowledge

Fishers have developed local or TEK of the resources that they depend on based on their daily interactions with these resources over long time periods (Berkes 1999). Fishers thus have a great deal to offer scientists and managers in terms of holistic understanding of marine ecosystem dynamics in the areas they know well (Johannes 1978, 1998). TEK has been gathered, synthesized, and passed on orally, often in the form of anecdotes (Agrawal 1995; Johannes and Neis 2007). TEK is therefore context specific, untested, sometimes unreliable, and, until recently, very difficult for classically trained ecologists, oceanographers, or fisheries managers to accept, incorporate, or mesh with their own studies (Johannes and Neis 2007; Shackeroff and Campbell 2007). As a result, there exist far too many examples of marine research and management programs that have ignored fishers and their local knowledge (Johannes, Freeman, and Hamilton 2000). The consequences of ignoring fishers' knowledge include ignorant conclusions in stock assessments that have missed known seasonality or migration patterns or, worse, fisheries collapses. In Tarawa, Kiribati, for example, annual spawning runs

of bonefish were almost completely destroyed when causeways were built between islands surrounding the atoll that blocked seasonal spawning migrations. Based on interviews with older, experienced fishers from remote villages, a single relict spawning run was discovered and subsequently managed, leading to resurgence in the bonefish population (Johannes, Freeman, and Hamilton 2000). In another case highlighted in this Focus Section, marine protected area boundaries in Florida's Carysfort Reef were drawn to include a spawning aggregation but inadvertently excluded an adjacent but locally well-known black grouper spawning site (Gleason, Kellison, and Reid this issue). As will be developed in the following section, however, the divide between traditional ecological knowledge and that derived from scientists is not insurmountable (Agrawal 1995). Fishers' knowledge may be particularly valuable to scientists as we try to move to EBM (as noted in the preceding section), although scholars must be cognizant of delicate political, cultural, and power relationships and issues that arise in this type of research (Shackeroff and Campbell 2007). The time has come for putting fishers' knowledge *back* to work for conservation and marine resource management (Haggan, Neis, and Baird 2007).

We emphasize "back to work" because there are myriad examples of preindustrial societies effectively managing their fisheries based on TEK and local management structures, particularly in the Indo-Pacific (e.g., Johannes 1978; Berkes 1999). Unfortunately, however, many local traditional marine management systems are being abandoned or eclipsed by the onset and interaction with industrial fishing and modern centralized governance, often with negative effects on resources (e.g., McClanahan et al. 1997). Because local fishers' involvement in management is predicated on both their understanding of the complex systems in question, as well as their personal stake in the effective management of those resources, their involvement can lead to effective EBM solutions with high compliance. Modern examples are becoming more common (e.g., Drew 2005; Prigent et al. 2008). Indeed, most modern scholars and managers consider the involvement of fishers and their TEK as an integral component of effective fisheries management and ocean governance (National Research Council 1999, 2001;

Berkes, Colding, and Folke 2000; Berkes 2004; U.S. Commission on Ocean Policy 2004). Importantly for geographers, carefully planned and implemented studies of TEK and their integration with other traditional disciplinary studies are equally valuable and needed on land and in the sea.

### Stakeholder Involvement in and Governance of Common Property Resources

The world's oceans are the largest and most important of our common property resources. Their management has suffered the tragedy of the commons (Hardin 1968), and this has recently been attributed to insufficient governance by appropriate and effective institutions (Dietz, Ostrom, and Stern 2003). Crowder et al. (2006) indicate that declining marine ecosystem health is largely due to spatial and temporal mismatches between the scale of ecosystems and the jurisdiction of their management institutions (e.g., species that migrate across national boundaries). Their article articulated the need for large-scale ocean zoning based on "underlying topography, oceanography, and the distribution of biotic communities" (617). An increasing number of articles have argued that effective marine area governance can be predicated on the involvement of appropriate stakeholders, particularly fishers, in the process of adaptive management (e.g., St. Martin 2001; Christie et al. 2005; Christie and White 2007). Others have argued that marine systems can be viewed as complex and coupled social-ecological systems so their management should be addressed using a multidisciplinary approach that addresses the interactions among resources, resource users, and governance institutions (Mahon, McConney, and Roy 2008; McClanahan et al. 2008).

Comprehensive ocean zoning represents the convergence of inclusive governance and EBM and thus serves to mitigate conflicts in ocean use and to promote sustainability in fisheries and marine ecosystems (Halpern et al. 2008; Norse 2010). Ocean zoning is implicitly spatial. St. Martin and Hall-Arber (2008b) suggested that whereas many physical properties of marine systems are increasingly well expressed in marine GIS systems for planning, community resource use is less well represented. Recent

studies seek to fill this gap by offering methods to work with fishers to map their resource use in GIS layers that can be considered as an integral component in marine spatial planning (Lunn and Dearden 2006; St. Martin and Hall-Arber 2008a, 2008b). Planning and management of the Great Barrier Reef Marine Park, for example, included comprehensive stakeholder involvement including GIS data layers showing resource use (Day 2002). There are a growing number of examples of the comanagement of common property coastal and marine resources that illustrate the concepts in this section. For example, community-based cooperatives contribute to the sustainable management of the Sian Ka'an Biosphere Reserve in Mexico. As part of their involvement, they manage cooperatively a sustainable lobster fishery through good governance, transparency, and solid science (Sosa-Cordero, Liceaga-Correa and Seijo 2008). Integrated coastal management programs have also seen some success in the Philippines (White, Deguit, and Jatulan 2006) and Trinidad and Tobago (Tompkins, Adger, and Brown 2002). These studies provide local examples addressing commons management but the issue remains as one of the world's greatest challenges (Ostrom et al. 1999).

### Synthesis of the Focus Section

Although this Focus Section is far from comprehensive, this lead article and the articles herein provide foundational data and holistic approaches to address declining health and resilience of ocean resources. We argue for the expansion of marine ecosystem-based spatial management based on integrated multidisciplinary studies of underlying geomorphology as a proxy for marine habitats, studies of marine connectivity, fishers' TEK, the critical analysis of institutions and governance, and the broad involvement of stakeholders in the entire process.

The articles in this section focus on the ecology and management of tangible and specific subset of ocean governance and management issues and areas—the ecology and management of reef fish breeding areas. We recommend extensive documentation of the timing and location of multispecies spawning aggregation areas starting with geomorphology and fisher interviews as primary sources of information. The

protection of these critical breeding and feeding habitats, through cooperative management, within seascape networks of reserves will contribute to regional reef ecosystem resilience.

In summary, the five articles collected here illustrate the following:

1. State-of-the-art examples of how researchers have classified, integrated, and analyzed physical and ecological data sources to reveal geomorphology as a proxy for marine habitat, specifically, reef fish spawning aggregations (Gleason, Kellison, and Reid this issue; Heyman this issue; Coleman, Scanlon, and Koenig this issue)
2. How spatial models of larval transport can illustrate the connectivity between spawning areas and nursery grounds via ocean currents (Fischer et al. this issue)
3. How these scientific results, along with the active participation of stakeholders (such as fishers and their TEK), can be effectively used in the design of functional marine reserve networks; hence a demonstration of marine EBM in action (Heyman this issue)

These articles provide compelling examples of the important role of geographic inquiry and its applications in marine environments. As a discipline, geography is eclectic and multidisciplinary, yet holistic and integrative. This Focus Section endeavors to provide that breadth (marine geomorphology, marine ecology, marine spatial planning based on ecosystem principles, stakeholder participation in governance of common property resources, and the design of functional marine reserve networks) along with depth of inquiry into a representative set of case studies that illustrates the breadth but focuses on largely one issue—marine spatial planning that aims to conserve vulnerable and valuable reef fish spawning aggregations and their connectivity, based on studies of geomorphology as a proxy for habitat and modeling studies of marine connectivity and through the involvement and participation of fishers and their TEK. ■

## Literature Cited

- Agrawal, A. 1995. Dismantling the divide between indigenous and scientific knowledge. *Development and Change* 26:413–39.
- Allison, G. W., S. D. Gaines, J. Lubchenco, and H. P. Possingham. 2003. Ensuring persistence of marine reserves: Catastrophes require adopting an insurance factor. *Ecological Applications* 13 (1, Suppl.): S8–S24.
- Andréfouët, S., M. J. Costello, D. P. Faith, S. Ferrier, G. N. Geller, R. Höft, N. Jürgens, et al. 2008. *The GEO Biodiversity Observation Network concept document*. GEO—Group on Earth Observations, Geneva, Switzerland.
- Aswani, S., and M. Lauer. 2006. Benthic mapping using local aerial photo interpretation and resident taxa inventories for designing marine protected areas. *Environmental Conservation* 33:263–73.
- Ball, I. R., and H. P. Possingham. 2000. MARXAN v1.2, marine reserve design using spatially explicit annealing, a manual. [http://www.uq.edu.au/marxan/docs/marxan\\_manual.1.8.2.pdf](http://www.uq.edu.au/marxan/docs/marxan_manual.1.8.2.pdf) (last accessed 17 December 2008).
- Balmford, A., P. Gravestock, N. Hockley, C. J. Mclean, and C. M. Roberts. 2004. The worldwide costs of marine protected areas. *Proceedings of the National Academy of Sciences* 101 (26): 9694–97.
- Beck, M. W., K. L. Heck Jr., K. W. Able, D. L. Childers, D. B. Eggleston, B. M. Gillanders, B. Halpern, et al. 2001. The identification, conservation, and management of estuarine and marine fish nurseries for fish and invertebrates. *Bioscience* 51:633–41.
- Berkeley, S. A., M. A. Hixon, N. J. Larson, and M. S. Love. 2004. Fisheries sustainability via protection of age structure and spatial distribution of fish populations. *Fisheries* 29 (8): 23–32.
- Berkes, F. 1999. *Sacred ecology: Traditional ecological knowledge and resource management*. London and New York: Taylor & Francis.
- Berkes, F. 2004. Rethinking community based conservation. *Conservation Biology* 18 (3): 621–30.
- Berkes, F., J. Colding, and C. Folke. 2000. Rediscovery of traditional ecological knowledge as adaptive management. *Ecological Applications* 10:1251–62.
- Burke, L., and J. Maidens. 2004. *Reefs and risk in the Caribbean*. Washington, DC: World Resources Institute.
- Burne, R. V., and C. A. Parvey. 2002. Marine geography and the benthic habitat: Domains of the Australian Ocean Territory. In *Marine geography: GIS for the oceans and seas*, ed. J. Breman, 127–36. Redlands, CA: ESRI Press.
- Christie, P., D. L. Fluharty, A. T. White, L. Eisma-Osorio, and W. Jatulan. 2007. Assessing the feasibility of ecosystem-based fisheries management in tropical contexts. *Marine Policy* 31:239–50.
- Christie, P., K. Lowry, A. T. White, E. G. Oracion, L. Sievanen, R. S. Pomeroy, R. B., Pollnac, J. M. Patlis, and R.-L. V. Eisma. 2005. Key findings from a multidisciplinary examination

- of integrated coastal management process sustainability. *Ocean & Coastal Management* 48:468–83.
- Christie, P., and A. T. White. 2007. Best practices for improved governance of coral reef marine protected areas. *Coral Reefs* 26:1047–56.
- Claro, R., and K. Lindeman. 2003. Spawning aggregation sites of snapper and grouper species (Lutjanidae and Serranidae) on the insular shelf of Cuba. *Gulf and Caribbean Research Institute* 14:91–106.
- Claydon, J. 2004. Spawning aggregations of coral reef fishes: Characteristics, hypotheses, threats, and management. *Oceanography and Marine Biology: An Annual Review* 42:265–302.
- Coleman, F. C., W. F. Figueira, J. Ueland, and L. B. Crowder. 2004. The impact of United States recreational fisheries on marine fish populations. *Science* 305:1958–60.
- Costanza, R., R. d'Arge, R. de Groot, S. Farber, M. Grasso, B. Hannon, K. Limburg, et al. 1997. The value of the world's ecosystem services and natural capital. *Nature* 387:253–60.
- Costello, M. 2009. Distinguishing marine habitat classification concepts for ecological data management. *Marine Ecology Progress Series* 397:253–68.
- Creel, L. 2003. Ripple effects: Population and coastal regions. Washington, DC: Population Reference Bureau. <http://www.prb.org/Publications/PolicyBriefs/RippleEffectsPopulationandCoastalRegions.aspx> (last accessed 1 June 2011).
- Crowder, L. B., E. L. Hazen, N. Avissar, R. Bjorkland, C. Latanich, and M. B. Ogburn. 2008. The impacts of fisheries on marine ecosystems and the transition to ecosystem-based management. *Annual Review of Ecology Evolution and Systematics* 39:259–78.
- Crowder, L. B., G. Osherenko, O. R. Young, S. Airamé, E. A. Norse, N. Baron, J. C. Day, et al. 2006. Resolving mismatches in U.S. ocean governance. *Science* 313:617–18.
- Day, J. 2002. Zoning—Lessons from the Great Barrier Reef Marine Park. *Ocean and Coastal Management* 45:139–56.
- Dietz, T., E. Ostrom, and P. C. Stern. 2003. The struggle to govern the commons. *Science* 302:1907–12.
- Domeier, M. L., and P. L. Colin. 1997. Tropical reef fish spawning aggregations: Defined and reviewed. *Bulletin of Marine Science* 60 (3): 698–726.
- Drew, J. A. 2005. Use of traditional ecological knowledge in marine conservation. *Conservation Biology* 19:1286–93.
- Erdey-Heydorn, M. D. 2008. An ArcGIS seabed characterization toolbox developed for investigating benthic habitats. *Marine Geodesy* 31:318–58.
- Food and Agriculture Organization of the United Nations (FAO). 2004. *The state of world fisheries and aquaculture*. Rome, Italy: FAO.
- . 2009. *The state of world fisheries and aquaculture 2008*. Rome, Italy: FAO.
- Green, E. P., P. J. Mumby, A. J. Edwards, and C. D. Clark. 1996. A review of remote sensing for the assessment and management of tropical coastal resources. *Coastal Management* 24:1–40.
- Grober-Dunsmore, R., T. K. Frazer, J. P. Beets, W. J. Lindberg, P. Zwick, and N. A. Funicelli. 2008. Influence of landscape structure on reef fish assemblages. *Landscape Ecology* 23:37–53.
- Grober-Dunsmore, R., and B. D. Keller, eds. 2008. *Caribbean connectivity: Implications for marine protected area management*. Marine Sanctuaries Conservation Series ONMS-08-07. Silver Spring, MD: National Oceanic and Atmospheric Administration. <http://sanctuaries.noaa.gov/science/conservation/carib.html> (last accessed 1 June 2011).
- Haggan, N., B. Neis, and I. G. Baird. 2007. Putting fishers' knowledge to work. In *Fishers' knowledge in fisheries science and management*, ed. N. Haggan, B. Neis, and G. Baird, 35–40. Coastal Management Sourcebooks series. Paris: UNESCO.
- Halpern, B. S. 2003. The impact of marine reserves: Do reserves work and does reserve size matter? *Ecological Applications* 13 (1): S117–S137.
- Halpern, B. S., K. L. McLeod, A. A. Rosenberg, and L. B. Crowder. 2008. Managing for cumulative impacts in ecosystem-based management through ocean zoning. *Ocean & Coastal Management* 51 (3): 203–11.
- Halpern, B. S., and R. R. Warner. 2003. Matching marine reserve design to reserve objectives. *Proceedings of the Royal Society of London B* 270 (1527): 1871–78.
- Hardin, G. 1968. Tragedy of the commons. *Science* 162:1243–48.
- Hastings, A., and W. W. Botsford. 2003. Comparing designs of marine reserves for fisheries and for biodiversity. *Ecological Applications* 13 (1): S65–S70.
- Hayden, B. P., G. C. Ray, and R. Dolan. 1984. Classification of coastal and marine environments. *Environmental Conservation* 11:199–207.
- Heap, A. D., and P. T. Harris. 2008. Geomorphology of the Australian margin and adjacent seafloor. *Australian Journal of Earth Sciences* 55 (4): 555–85.
- Hedgpeth, J. W. 1957a. Classification of marine environments. *Geological Society of America Memoirs* 67:17–28.
- . 1957b. Marine biogeography. *Geological Society of America Memoirs* 67:359–82.
- Heyman, W. D., and B. Kjerfve. 2008. Characterization of transient multi-species reef fish spawning aggregations at Gladden Spit, Belize. *Bulletin of Marine Science* 83 (3): 531–51.

- Hoegh-Guldberg, O., P. J. Mumby, A. J. Hooten, R. S. Steneck, P. Greenfield, E. Gomez, D. R. Harvell, et al. 2007. The carbon crisis: Coral reefs under rapid climate change and ocean acidification. *Science* 318:1737–42.
- Iampietro, P. J., M. A. Young, and R. G. Kvittek. 2008. Multivariate prediction of rockfish habitat suitability in Cordell Bank National Marine Sanctuary and Del Monte Shalebeds, California, USA. *Marine Geodesy* 31:359–71.
- Jackson, J. B. C., M. X. Kirby, W. H. Berger, K. A. Bjorndal, L. W. Botsford, B. J. Bourque, R. H. Bradbury, et al. 2001. Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293 (5530): 629–37.
- Johannes, R. E. 1978. Reproductive strategies of coastal marine fishes in the tropics. *Environmental Biology of Fishes* 3:65–84.
- . 1998. The case for data-less marine resource management: Examples from tropical nearshore fisheries. *Trends in Ecology & Evolution* 13:243–46.
- Johannes, R. E., M. M. R. Freeman, and R. J. Hamilton. 2000. Ignore fishers' knowledge and miss the boat. *Fish and Fisheries* 1:257–71.
- Johannes, R. E., and B. Neis. 2007. The value of anecdote. In *Fishers' knowledge in fisheries science and management*, ed. N. Haggan, B. Neis, and G. Baird, 41–58. Coastal Management Sourcebooks series. Paris: UNESCO.
- Kendall, M. S., and T. Miller. 2008. The influence of thematic and spatial resolution on maps of a coral reef ecosystem. *Marine Geodesy* 31:75–102.
- Klein, C. J., C. Steinback, A. J. Scholz, and H. P. Possingham. 2008. Effectiveness of marine reserve networks in representing biodiversity and minimizing impact to fishermen: A comparison of two approaches used in California. *Conservation Letters* 1 (1): 44–51.
- Kobara, S., and W. D. Heyman. 2008. Geomorphometric patterns of Nassau grouper (*Epinephelus striatus*) spawning aggregation sites in the Cayman Islands. *Marine Geodesy* 31 (4):231–45.
- . 2010. Sea bottom morphology of multi-species spawning sites in Belize. *Marine Ecology Progress Series* 31:231–45.
- Koenig, C. C., F. C. Coleman, L. A. Collins, Y. Sadovy, and P. L. Colin. 1996. Reproduction of gag (*Mycteroperca microlepis*) (Pisces: Serranidae) in the eastern Gulf of Mexico and the consequences of fishing spawning aggregations. In *ICLARM Conference 48, ICLARM Conference Proceeding*, 307–23. Manila, Philippines: International Center for Living Aquatic Resources Management.
- Koenig, C. C., F. C. Coleman, C. B. Grimes, G. R. Fitzhugh, K. M. Scanlon, C. T. Gledhill, and M. Grace. 2000. Protection of fish spawning habitat for the conservation of warm-temperate reef-fish fisheries of shelf-edge reefs of Florida. *Bulletin of Marine Science* 66 (3): 593–616.
- Kracker, L., M. Kendall, and G. McFall. 2008. Benthic features as a determinant for fish biomass in Gray's Reef National Marine Sanctuary. *Marine Geodesy* 31:267–80.
- Lanier, A., C. Romsos, and C. Goldfinger. 2007. Seafloor habitat mapping on the Oregon continental margin: A spatially nested GIS approach to mapping scale, mapping methods, and accuracy quantification. *Marine Geodesy* 30:51–76.
- Leis, J. M. 1987. Review of the early life history of tropical groupers (Serranidae) and snappers (Lutjanidae). In *Tropical snappers and groupers*, ed. J. J. Polovina and S. Ralston, 189–238. Boulder, CO: Westview.
- Leslie, H., M. Ruckelshaus, I. R. Ball, S. Andelman, and H. P. Possingham. 2003. Using siting algorithms in the design of marine reserve networks. *Ecological Applications* 13 (1): S185–S198.
- Levin, S. A., and J. Lubchenco. 2008. Resilience, robustness and marine ecosystem-based management. Corvallis, OR: *BioScience* 58:27–32.
- Lunn, K. E., and P. Dearden. 2006. Fishers' needs in marine protected area zoning: A case study from Thailand. *Coastal Management* 34:183–98.
- Mahon, R., P. McConney, and R. N. Roy. 2008. Governing fisheries as complex adaptive systems. *Marine Policy* 32:104–12.
- McClanahan, T. R., J. C. Castilla, A. T. White, and O. Defeo. 2008. Healing small-scale fisheries by facilitating complex socio-ecological systems. *Reviews in Fisheries Biology and Fisheries* 19:33–47.
- McClanahan, T. R., H. Glaesel, J. Rubens, and R. Kiambo. 1997. Traditional fisheries management, its decay, and effect on fisheries yields and the coral reef ecosystems of Southern Kenya. *Environmental Conservation* 24 (2): 105–20.
- McLeod, K. L., J. Lubchenco, S. R. Palumbi, and A. A. Rosenberg. 2005. *Scientific consensus statement on marine ecosystem-based management*. Corvallis, OR: Communication Partnership for Science and the Sea. [http://www.compassonline.org/sites/all/files/document\\_files/EBM.Consensus.Statement.v12.pdf](http://www.compassonline.org/sites/all/files/document_files/EBM.Consensus.Statement.v12.pdf) (last accessed 1 June 2011).
- Millennium Ecosystem Assessment. 2005. *Living beyond our means: Natural assets and human well being*. Washington, DC: World Resources Institute.
- Mumby, P. J., K. Broad, D. R. Brumbaugh, C. P. Dahlgren, A. R. Harborne, A. Hastings, K. E. Holmes, C. V. Kappel, F. Micheli, and J. N. Sanchiro. 2008. Coral reef habitats as surrogates of species, ecological functions, and ecosystem services. *Conservation Biology* 22 (4): 941–51.
- Myers, R. A., and B. Worm. 2003. Rapid worldwide depletion of predatory fish communities. *Nature* 423:280–83.

- National Research Council. 1999. *Sustaining marine fisheries*. Washington, DC: National Academy Press.
- . 2001. *Marine protected areas: Tools for sustaining ocean ecosystem*. Washington, DC: National Academies Press.
- NatureServe. 2008. Ecosystem-based management tools network. <http://www.ebmttools.org> (last accessed 17 December 2008).
- Norse, E. A. 2010. Ecosystem-based spatial planning and management of marine fisheries: Why and how? *Bulletin of Marine Science* 86 (2): 179–95.
- Okubo, A., and S. Levin. 2002. The basics of diffusion. In *Diffusion and ecological problems: Modern perspectives*. 2nd ed., ed. A. Okubo and S. Levin, 10–30. New York: Springer.
- Ostrom, E., J. Burger, C. B. Field, R. B. Norgaard, and D. Policansky. 1999. Revisiting the commons: Local lessons, global challenges. *Science* 284:278–82.
- Palumbi, S. R., P. A. Sandifer, J. D. Allan, M. W. Beck, D. G. Fautin, M. J. Fogarty, B. S. Halpern, et al. 2009. Managing for ocean biodiversity to sustain marine ecosystem services. *Frontiers in Ecology and the Environment* 4:204–11.
- Pauly, D., V. Christensen, J. Dalsgaard, R. Froese, and F. C. Torres Jr. 1998. Fishing down marine food webs. *Science* 279:860–63.
- Pauly, D., V. Christensen, S. Guénette, T. J. Pitcher, U. R. Sumaila, C. J. Walters, R. Watson, & D. Zeller. 2002. Towards sustainability in world fisheries. *Nature* 418:689–95.
- Peterson, C. H., H. C. Summerson, E. Thomson, H. S. Lenihan, J. Grabowski, L. Manning, F. Micheli, and G. Johnson. 2000. Synthesis of linkages between benthic and fish communities as a key to protecting essential fish habitat. *Bulletin of Marine Science* 66 (3): 759–74.
- Pikitch, E. K., C. Santora, E. A. Babcock, A. Bakun, R. Bonfil, D. O. Conover, P. Dayton, et al. 2004. Ecosystem-based fishery management. *Science* 305:346–47.
- Pittman, S. J., C. Caldwell, and S. D. Hile. 2007. Using seascape types to explain the spatial patterns of fish in the mangroves of SW Puerto Rico. *Marine Ecology Progress Series* 348:273–84.
- Pittman, S. J., C. A. McAlpine, and K. M. Pittman. 2004. Linking fish and prawns to their environment: A hierarchical landscape approach. *Marine Ecology Progress Series* 283:233–254.
- Possingham, H., I. Ball, and S. Andelman. 2000. Mathematical methods for identifying representative reserve networks. In *Quantitative methods for conservation biology*, ed. S. Ferson and M. Burgman, 291–305. Berlin: Springer-Verlag.
- Prigent, M., G. Fontenelle, M.-J. Rochet, and V. M. Trenkel. 2008. Using cognitive maps to investigate fishers' ecosystem objectives and knowledge. *Ocean & Coastal Management* 51:450–62.
- Psuty, N. P., P. E. Steinberg, and D. J. Wright. 2004. Coastal and marine geography. In *Geography in America at the dawn of the 21st century*, ed. G. L. Gaile and C. J. Willmott, 314–25. New York: Oxford University Press.
- Roberts, C. M. 1997. Connectivity and management of Caribbean coral reefs. *Science* 278:1454–57.
- Roberts, C. M., J. A. Bohnsack, F. Gell, J. P. Hawkins, and R. Goodridge. 2001. Effects of marine reserves on adjacent fisheries. *Science* 294:1920–23.
- Roff, J. C., and M. E. Taylor. 2000. National frameworks for marine conservation—A hierarchical geophysical approach. *Aquatic Conservation: Marine and Freshwater Ecosystems* 13:77–90.
- Roff, J. C., M. E. Taylor, and J. Laughren. 2003. Geophysical approaches to the classification, delineation and monitoring of marine habitats and their communities. *Aquatic Conservation: Marine and Freshwater Ecosystems* 13:77–90.
- Sadovy, Y., and M. Domeier. 2005. Are aggregation fisheries sustainable? Reef fish fisheries as a case study. *Coral Reefs* 24:254–62.
- Sadovy De Mitcheson, Y. S., A. Cornish, M. Domeier, P. L. Colin, M. Russell, and K. C. Lindeman. 2008. A global baseline for spawning aggregations of reef fishes. *Conservation Biology* 22 (5): 1233–44.
- Sala, E., R. Starr, and E. Ballesteros. 2001. Rapid decline of Nassau grouper spawning aggregations in Belize: Fishery management and conservation needs. *Fisheries* 26 (10): 23–30.
- Sandwell, D., S. Gille, J. A. Orcutt, and W. Smith. 2003. Bathymetry from space is now possible. *Eos, Transactions of the American Geophysical Union* 84 (5): 37, 44.
- Shackeroff, J. M., and L. M. Campbell. 2007. Traditional ecological knowledge in conservation research: Problems and prospects for their constructive engagement. *Conservation and Society* 5:343–60.
- Smith, M. D., C. A. Robbins, L. B. Crowder, B. S. Halpern, M. Turnipseed, J. L. Anderson, F. Asche, et al. 2010. Sustainability and global seafood. *Science* 327:784–86.
- Sosa-Cordero, E., M. L. A. Liceaga-Correa, and J. C. Seijo. 2008. The Punta Allen lobster fishery: Current status and recent trends. In *Case studies on fisheries self-governance*, ed. R. Townsend, R. Shotton, and H. Uchida, 149–62. FAO Fisheries Technical Paper No. 514. Rome, Italy: FAO.
- Spalding, M. D., H. E. Fox, G. R. Allen, N. Davidson, Z. A. Ferdaña, M. Finlayson, B. S. Halpern, et al. 2007. Marine ecoregions of the world: A bioregionalization of coastal and shelf areas. *Bioscience* 57 (7): 573–83.

- St. Martin, K. 2001. Making space for community resource management in fisheries. *Annals of the Association of American Geographers* 91:122–42.
- St. Martin, K., and M. Hall-Arber. 2008a. Creating a place for “community” in New England fisheries. *Human Ecology Review* 15:161–70.
- . 2008b. The missing layer: Geo-technologies, communities, and implications for marine spatial planning. *Marine Policy* 32:779–86.
- Steinberg, P. 1999. Navigating to multiple horizons: Toward a geography of open space. *The Professional Geographer* 51 (3): 366–75.
- Thresher, R. E. 1984. *Reproduction in reef fishes*. Neptune City, NJ: TFH Publications.
- Tompkins, E., W. N. Adger, and K. Brown. 2002. Institutional networks for inclusive coastal management in Trinidad and Tobago. *Environment and Planning A* 34:1095–1111.
- U.S. Commission on Ocean Policy. 2004. *An ocean blueprint for the 21st century*. Washington, DC: U.S. Commission on Ocean Policy.
- Warner, R. R., S. E. Swearer, and J. E. Caselle. 2000. Larval accumulation and retention: Implications for the design of marine reserves and essential fish habitat. *Bulletin of Marine Science* 66 (3): 821–30.
- Wedding, L. M., and A. M. Friedlander. 2008. Determining the influence of seascape structure on coral reef fishes in Hawaii using a geospatial approach. *Marine Geodesy* 31:246–66.
- Whyalen, L., C. V. Pattengill-Semmens, B. X. Semmens, P. G. Bush, and M. R. Boardman. 2004. Observations of a Nassau grouper, *Epinephelus striatus*, spawning aggregation site in Little Cayman, Cayman Islands, including multi-species spawning information. *Environmental Biology of Fishes* 70:305–13.
- White, A., E. Deguit, and W. Jatulan. 2006. Integrated coastal management in Philippine local governance: Evolution and benefits. *Coastal Management Journal* 34:287–302.
- Wilson, M. F. J., B. O’Connell, C. Brown, J. C. Guinan, and A. J. Grehan. 2007. Multiscale terrain analysis of multibeam bathymetry data for habitat mapping on the continental slope. *Marine Geodesy* 30:3–35.
- Worm, B., E. B. Barbier, N. Beaumont, J. E. Duffy, C. Folke, B. S. Halpern, J. B. C. Jackson, et al. 2006. Impacts of biodiversity loss on ocean ecosystems services. *Science* 314:787–90.
- Worm, B., R. Hilborn, J. K. Baum, T. A. Branch, J. S. Collie, C. Costello, M. J. Fogarty, et al. 2009. Rebuilding global fisheries. *Science* 325:578–85.
- Wright, D. J. 1999. Getting to the bottom of it: Tools, techniques, and discoveries of deep ocean geography. *The Professional Geographer* 51:426–39.
- . 2003. Introduction. In *Undersea with GIS*, ed. D. J. Wright, xiii–xvi. Redlands, CA: ESRI Press.
- Wright, D. J., B. Donahue, and D. F. Naar. 2002. Seafloor mapping and GIS coordination at America’s remotest national marine sanctuary (American Samoa). In *Undersea with GIS*, ed. D. J. Wright, 33–63. Redlands, CA: ESRI Press.
- Wright, D. J., and W. D. Heyman. 2008a. Marine and coastal GIS for geomorphology, habitat mapping, and marine reserves. *Marine Geodesy* 31 (4): 1–8.
- . 2008b. Marine geomorphology as a determinant for essential life habitat: An ecosystem management approach to planning for marine reserve networks. <http://marinecoastalgis.net/aag08> (last accessed 17 December 2008).
- Zacharias, M. A., D. E. Howes, J. R. Harper, and P. Wainwright. 1998. The British Columbia marine ecosystem classification: Rationale, development, and verification. *Coastal Management* 26:105–24.
- Zacharias, M. A., and J. C. Roff. 2000. A hierarchical ecological approach to conserving marine biodiversity. *Conservation Biology* 14 (5): 1327–34.

WILLIAM D. HEYMAN is an Associate Professor in the Department of Geography at Texas A&M University, College Station, TX 77845. E-mail: [whyeman@tamu.edu](mailto:whyeman@tamu.edu). His research entails physical, biological, and social aspects of marine and coastal conservation and management.

DAWN J. WRIGHT is a Professor of Geography and Oceanography at Oregon State University, Corvallis, OR 97331–5506. E-mail: [dawn@dusk.geo.orst.edu](mailto:dawn@dusk.geo.orst.edu). Her research interests include GIScience, ocean informatics, benthic terrain and habitat characterization, and the processing and interpretation of high-resolution bathymetry and underwater imagery.